

Telenor Font Family

NEWLYN

The Telenor typeface has been developed by Miles Newlyn and is based on the work of Magnus Rakeng. It is the proprietary typeface of the Telenor company. Developed in 2005, it is a complete redraw of the earlier typeface, and has a broad character set that includes Cyrillic.

Miles Newlyn is an internationally respected designer, having created identities for some of the world largest organisations, including Honda and Unilever. Newlyn is also an expert in corporate typography, designing typefaces for clients such as Citibank, British Telecom, and the Tate Gallery.

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION

16/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to procure,

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individuals of each species are born

than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend to propagate its new and modified

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus</i> sp.	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as each man tries to

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individ-

uals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
Apereiodon affinis	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
Apereiodon piracicabae	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
Hypostomus ragani	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
Hypostomus sp.	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
Prochilodus lineatus	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
Steindachnerina corumbae	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
Steindachnerina insculpta	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS
NATURALIST, I WAS MUCH STRUCK
WITH CERTAIN FACTS IN THE DISTRI-

16/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many more individ-

uals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRI-

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as each

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As

many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
Apereiodon affinis	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
Apereiodon piracicabae	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
Hypostomus ragani	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
Hypostomus sp.	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
Prochilodus lineatus	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
Steindachnerina corumbae	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
Steindachnerina insculpta	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRI-

16/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As many

more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at work, as each

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As

many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
Apereiodon affinis	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
Apereiodon piracicabae	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
Hypostomus ragani	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
Hypostomus sp.	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
Prochilodus lineatus	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
Steindachnerina corumbae	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
Steindachnerina insculpta	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN

16/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. As

many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN

16/18pt

When the first tendency was once displayed, methodical selection *and the inherited effects of compulsory training in each successive generation* would soon complete the work; and unconscious selection is still at

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of

our greatest philosophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving,

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN

16/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of

our greatest philosophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving,

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insculpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

26/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN

16/18pt

When the flrst tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection

8/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one

of our greatest philosophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better

5/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Apereiodon affinis</i>	1	"caniveteš	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apereiodon piracicabae</i>	6	"caniveteš	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudoš	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudoš	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbataš	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiruš	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina inculpta</i>	47	"saguiruš	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

Telenor

- Deutsch Als ich an Bord des Königlichen Schiffs Beagle als Naturforscher Südamerika erreichte, ward ich überrascht von der Wahrnehmung gewisser Thatsachen in der Vertheilung der Bewohner und in den geologischen Bez-
- Español Cuando estaba como naturalista a bordo del Beagle, buque de la marina real, me impresionaron mucho ciertos hechos que se presentan en la distribución geográfica de los seres orgánicos que viven en América del Sur y en las
- Français Les rapports géologiques qui existent entre la faune actuelle et la faune éteinte de l'Amérique méridionale, ainsi que certains faits relatifs à la distribution des êtres organisés qui peuplent ce continent, m'ont profondément
- Italiano Io mi trovavo a bordo del vascello di S.M.Britannica The Beagle nella qualità di naturalista, allorchè fui vivamente colpito da certi fatti nella distribuzione degli esseri organizzati che popolano l'America meridionale e dai rap-
- Suomi Kun katsomme jalostamiemme kasvien ja eläinten lajeja ja alalajeja yksilöinä, ensimmäinen silmään pistävä asia on, että ne yleisesti ottaen eroavat toisistaan huomattavasti enemmän kuin kuin minkään luonnonvaraiset lajin

Telenor

Dansk Darwin's tur i H.M.S. Beagle startede i april 1838 og sluttede i oktober 1838. Han var nok den mest banebrydende tænker i den moderne biologi. Under turen i skibet beagle var han i land i mange forskellige lande og under nogle af

Česky Po svém návratu z plavby rozvinul svou teorii přirozeného výběru. Věděl, že přírodovědci mezi duchovními, ač jeho přátelé, jsou přesvědčeni, že je to bestiální kacířství, podkopávající zázračná ospravedlnění pro společnost a věděl,

Türkçe HMS Beagle'in yolculuğu iki yerine beş yıl sürdü. Darwin, yolculuk boyunca çok çeşitli jeolojik oluşumlar, fosiller ve canlılar keşfetti, ve bunlardan örnekler topladı. Fırsat buldukça Cambridge'e keşiflerini anlatan ayrıntılı me-

Română Charles Darwin își amintește despre tatăl său ca fiind "cel mai înțelept dintre oameni". De asemenea, Darwin îl considera ca fiind unul dintre cei mai buni cunoscători ai sufletului omenesc. Cu toate acestea, este greu de înțeles

Português A viagem do Beagle durou quatro anos e nove meses, dois terços dos quais Darwin esteve em terra firme. Ele estudou uma rica variedade de características geológicas, fósseis, organismos vivos e conheceu muitas pessoas, entre

Telenor

Polski Niewiele brakowało, by Darwin w ogóle nie znalazł się na Beagle. Gdy w czasie letnich ferii 1838 roku przebywał z profesorem Sedgwickiem na wycieczce geologicznej w Walii, do jego mieszkania nadeszły dwa ważne listy, jeden

Cymraeg Yn ystod ei deithiau ar y Beagle astudiodd Darwin ddaear-eg cyfandiroedd ac ynysoedd yn ogystal â nifer o anifeiliaid, planhigion a ffosiliau. Mewn ffordd drefnus iawn, casglodd nifer enfawr o enghreifftiau nad oedd neb wedi eu

Nederlands Aan boord van de H.M.S. Beagle werd ik, als natuuronderzoeker, sterk getroffen door bepaalde feiten omtrent de verspreiding van de fauna en flora van Zuid-Amerika, en de geologische relaties tussen de tegenwoordige en de