

Luvbug Font Family

NEWLYN

Luvbug is the direct descendant of the modular lettering system designed for liquid crystal display.

Fused, flattened, outlined and individualized, then inflated and turbo pimped, a flagship for happiness and optimism; the Cooper Black of post-millennium delirium.

Luvbug Outline provides a subtle twist on the interplay between positive and negative forms; strokes and counters.

Luvbug is suitable for a variety of applications including music, entertainment and packaging.

Luvbug has been created by Miles Newlyn, an internationally respected designer, having created identities for some of the world largest organisations, including Honda and Unilever. Newlyn is also an expert in corporate typography, designing typefaces for clients such as Citibank, British Telecom, and the Tate Gallery.

32/28pt

**WHEN ON BOARD H.M.S. BEAGLE,
AS NATURALIST, I WAS MUCH
STRUCK WITH CERTAIN FACTS IN**

18/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still

9/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our

greatest philosophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of sur-

6/8pt

Species	C	Common Name	n	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apeceiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apeceiodon picacicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus raganii</i>	22	"caseudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"caseudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"cucimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insepulta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

32/28pt

**WHEN ON BOARD H.M.S. BEAGLE,
AS NATURALIST, I WAS MUCH
STRUCK WITH CERTAIN FACTS IN**

18/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still

9/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our

greatest philosophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance

6/8pt

Species	C	Common Name	n	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 – DETRITIVORES														
<i>Apeceiodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Apeceiodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"caseudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"caseudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina corumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina insepulta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

32/28pt

**WHEN ON BOARD H.M.S. BEAGLE,
AS NATURALIST, I WAS MUCH
STRUCK WITH CERTAIN FACTS IN**

18/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still

9/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geological relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of

our greatest philosophers. As many more individuals of each species are born than can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of sur-

6/8pt

Species	C	Common Name	N	1	2	3	4	5	6	7	8	9	10	11
GROUP 01 — DETRITIVORES														
<i>Aperciodon affinis</i>	1	"canivete"	169	0.03	10.16	—	0.01	0.04	0.01	—	17.24	—	72.32	72.32
<i>Aperciodon piracicabae</i>	6	"canivete"	69	0.04	6.66	—	0.05	—	—	—	11.34	1.52	0.74	79.65
<i>Hypostomus ragani</i>	22	"cascudo"	111	—	0.30	0.19	—	—	—	—	3.36	0.01	0.49	95.64
<i>Hypostomus sp.</i>	23	"cascudo"	19	0.01	0.15	—	—	—	—	—	9.50	1.45	—	88.88
<i>Prochilodus lineatus</i>	41	"curimbata"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.58	96.28
<i>Steindachnerina eorumbae</i>	46	"saguiru"	289	0.07	0.43	0.04	0.01	—	—	—	3.65	0.01	0.48	95.32
<i>Steindachnerina inseulpta</i>	47	"saguiru"	395	0.02	1.22	0.07	0.01	—	—	—	1.93	—	96.62	96.62

32/28pt

WHEN ON BOARD H.M.S. BEAGLE, AS NATURALIST, I WAS MUCH STRUCK WITH CERTAIN FACTS IN THE DISTRIBUTION OF

18/18pt

When the first tendency was once displayed, methodical selection and the inherited effects of compulsory training in each successive generation would soon complete the work; and unconscious selection is still at work, as each man tries to procure,

9/10pt

When on board H.M.S. Beagle, as naturalist, I was much struck with certain facts in the distribution of the inhabitants of South America, and in the geographical relations of the present to the past inhabitants of that continent. These facts seemed to me to throw some light on the origin of species—that mystery of mysteries, as it has been called by one of our greatest philosophers. In many more individuals of each species are born than

can possibly survive; and as, consequently, there is a frequently recurring struggle for existence, it follows that any being, if it vary however slightly in any manner profitable to itself, under the complex and sometimes varying conditions of life, will have a better chance of surviving, and thus be naturally selected. From the strong principle of inheritance, any selected variety will tend to propagate its new and modified form.

6/8pt

Species	C	Common Name	0	1	2	3	4	5	6	7	8	9	10	11
GROUP 01—DETMANNORES														
<i>Operculodon affinis</i>	0	"canine"	150	0.03	0.15	—	0.01	0.04	0.01	—	1725	—	72.52	72.52
<i>Operculodon phocaenoides</i>	6	"canine"	50	0.06	6.66	—	0.05	—	—	—	11.50	1.52	0.76	76.65
<i>Hypostomus agassizii</i>	22	"cassid"	111	—	0.20	0.19	—	—	—	—	2.22	0.01	0.49	95.65
<i>Hypostomus sp.</i>	23	"cassid"	10	0.01	0.15	—	—	—	—	—	0.50	0.05	—	66.66
<i>Pseudorasbora lineatus</i>	01	"cassid"	150	—	0.02	—	0.07	—	—	—	2.05	—	1.50	96.22
<i>Steindachnerina corembae</i>	05	"cassid"	220	0.07	0.05	0.04	0.01	—	—	—	2.55	0.01	0.00	95.22
<i>Steindachnerina lineolatus</i>	07	"cassid"	205	0.02	1.22	0.07	0.01	—	—	—	1.02	—	0.62	96.62

Luvbug

Deutsch

Als ich an Bord des Königlichen Schiffs Beagle als Naturforscher Südamerika erreichte, ward ich überrascht von der Wahrnehmung gewisser Thatsachen in der Vertheilung der Bewohner und in den geologischen Beziehungen zwischen

Español

Cuando estaba como naturalista a bordo del Beagle, buque de la marina real, me impresionaron mucho ciertos hechos que se presentan en la distribución geográfica de los seres orgánicos que viven en América del Sur y en las relaciones

Français

Les rapports géologiques qui existent entre la faune actuelle et la faune éteinte de l'Amérique méridionale, ainsi que certains faits relatifs à la distribution des êtres organisés qui peuplent ce continent, m'ont profondément frappé

Italiano

Io mi trovavo a bordo del vascello di S.M.Britannica The Beagle nella qualità di naturalista, allorchè fui vivamente colpito da certi fatti nella distribuzione degli esseri organizzati che popolano l'America meridionale e dai rapporti

Nederlands

Aan boord van de H.M.S. Beagle werd ik, als natuuronderzoeker, sterk getroffen door bepaalde feiten omtrent de verspreiding van de fauna en flora van Zuid-Amerika, en de geologische relaties tussen de tegenwoordige en de vroegere